

Global obstructions to gauge-invariance in chiral gauge theory on the lattice

David H. Adams

Math. dept. and Centre for the Subatomic Structure of Matter, University of
Adelaide, Adelaide, S.A. 5005, Australia.

email: dadams@maths.adelaide.edu.au

Abstract

It is shown that certain global obstructions to gauge-invariance in chiral gauge theory, described in the continuum by Alvarez-Gaumé and Ginsparg, are exactly reproduced on the lattice in the Overlap formulation at small non-zero lattice spacing (i.e. close to the classical continuum limit). As a consequence, the continuum anomaly cancellation condition $d_R^{abc} = 0$ is seen to be a necessary (although not necessarily sufficient) condition for anomaly cancellation on the lattice in the Overlap formulation.

1 Introduction

The Overlap formalism [1, 2] provides a potential solution to the important problem of constructing chiral gauge theories nonperturbatively on the lattice. Gauge anomalies are a central issue in this context: One would like to show that a gauge-invariant formulation of chiral gauge theories on the lattice is possible when the usual (continuum) anomaly cancellation conditions are satisfied. Conversely, when these conditions are not satisfied, one would like to see the continuum anomalies emerge in the lattice formulation. In particular, an interesting test for a lattice formulation of chiral gauge theory is whether it can capture the global obstructions to gauge-invariance of the continuum theory, which reflect the topological structure of the determinant line bundle over the gauge orbit space [3, 4].

In this paper we show that the overlap reproduces a basic class of such obstructions, described in the continuum by Alvarez-Gaumé and Ginsparg [3].¹ This is a further demonstration of the ability of the overlap to reproduce topological features of the continuum theory.² As a consequence we will see that the usual (i.e. continuum) local anomaly cancellation condition ((1.4) below) is a necessary condition for local anomaly-free chiral gauge theory on the lattice in the overlap formulation.

Global obstructions to the vanishing of local gauge anomalies in the overlap formulation were previously considered by H. Neuberger in [5]. An analogue of the the geometric perspective on the continuum chiral determinant as a section in a determinant line bundle was described for the overlap, and a class of global obstructions (which are naturally described in this setting) was explicitly constructed for the abelian theory. These were seen to vanish precisely when the fermion content of the theory satisfies

$$\sum_{\alpha} e_{\alpha}^3 = 0 \tag{1.1}$$

where the e_{α} 's label the irreducible U(1) representations of the fermion species. This is precisely the condition for cancellation of local gauge anomalies in the continuum theory. Thus, in the abelian case, the continuum anomaly cancellation condition (1.1) is a necessary condition for gauge-invariance of the lattice chiral determinant in the overlap formulation.³ Our result is a nonabelian variant of this.⁴

In the continuum theory, the global obstructions of Alvarez-Gaumé and Ginsparg arise as follows. Take spacetime to be the Euclidean 4-torus T^4 (the choice of di-

¹ The possibility that this could happen had been previously mentioned in [2].

² The parity-invariant overlap formulation of vector gauge theory in odd dimensions reproduces the global gauge anomaly of the continuum theory [6, 7], while for chiral gauge theory in even dimensions it has been shown to reproduce Witten's global anomaly [8, 9].

³ Herbert Neuberger has pointed out to me that these obstructions also arise for nonabelian gauge groups with U(1) subgroups, and that in four dimensions there are no additional restrictions in the nonabelian case, i.e. if all U(1) subgroups are free from anomalies then so is the nonabelian group.

⁴ Although the class of obstructions that we consider is different: the ones in [5] involve a torus in the orbit space of lattice gauge fields whereas ours involve a 2-sphere.

mension 4 is for concreteness; everything generalises to the T^{2n} case for arbitrary n), gauge group $SU(N)$, and consider a family ϕ_θ of gauge transformations parameterised by $\theta \in S^1$. If the fermion is in the fundamental representation then each ϕ_θ is a map from T^4 to $SU(N)$, and the family of these corresponds to a map $\Phi : T^5 \rightarrow SU(N)$, $\Phi(\theta, x) = \phi_\theta(x)$. The action of ϕ_θ on a gauge field A determines a family $\{A^\theta\}_{\theta \in S^1}$. The winding number of the phase of the chiral determinant around this circle-family of gauge fields is an obstruction to gauge-invariance of the chiral determinant (since if the determinant is gauge-invariant then it is constant around the family $\{A^\theta\}$ and the winding number vanishes). In [3] it was shown that this winding number equals the degree of the map Φ . Thus the obstruction is non-vanishing precisely when there exist maps $\Phi : T^5 \rightarrow SU(N)$ with non-vanishing degree (which happens, e.g., when $N = 3$).⁵

In the general case where the fermion content is specified by some arbitrary (typically reducible) representation R of $SU(N)$, the preceding generalises as follows. Instead of the degree of Φ , which is given by an expression of the form $d^{abc}h_{abc}$ where

$$d^{abc} = 2\text{tr}((T^a T^b + T^b T^a)T^c) \quad (1.2)$$

and the T^a 's are the generators of $SU(N)$, the obstruction is given by $d_R^{abc}h_{abc}$ where d_R^{abc} is given by (1.2) with T^a replaced by $R(T^a)$ etc. Using the well-known fact that there is a relation of the form⁶

$$d_R^{abc} = c(R)d^{abc} \quad (1.3)$$

⁵In [3] the spacetime was S^4 rather than T^4 , and a condition $\phi_0 \equiv 1$ was imposed, which allows Φ to be viewed as a map from S^5 to $SU(N)$. There is no essential difference with the present case though, since there is an isomorphism between the homotopy equivalence classes of $\text{Map}(S^k, SU(N))$ and $\text{Map}(T^k, SU(N))$.

⁶ The existence of a relation of this form can be seen as follows. Since the representation ring of $SU(N)$ is generated by the fundamental representation and its complex conjugate, it suffices to show (1.3) in the case where R is a tensor product of copies of the fundamental representation. Then $R(T^a) = T^a \otimes 1 \otimes \cdots \otimes 1 + 1 \otimes T^a \otimes 1 \otimes \cdots \otimes 1 + \dots + 1 \otimes \cdots \otimes 1 \otimes T^a$ etc, and it follows that $(R(T^a)R(T^b) + R(T^b)R(T^a))R(T^c) = (T^a T^b + T^b T^a)T^c \otimes 1 \otimes \cdots \otimes 1 + \dots + 1 \otimes \cdots \otimes 1 \otimes (T^a T^b + T^b T^a)T^c +$ terms which have a single T^a , T^b , or T^c in one of the tensor slots. Since $\text{tr}(T^a) = 0$ etc, it follows that the trace of these latter terms vanishes and we get (1.3).

we see that the obstruction in the general case is $c(R)$ times the degree of Φ . Thus in the case where $\text{Map}(T^5, SU(N))$ contains maps with non-vanishing degree a necessary condition for gauge-invariance of the chiral determinant is $c(R) = 0$, or

$$d_R^{abc} = 0 \tag{1.4}$$

Of course, this is just the usual (necessary and sufficient) condition for anomaly cancellation in the continuum theory (the non-abelian analogue of (1.1)).

In this paper we consider a lattice version of the preceding obstructions in the overlap formulation, and show that they reduce to the continuum obstructions in the classical continuum limit. Since the lattice and continuum obstructions are both specified by integers, it follows that the lattice obstruction is exactly equal to the continuum one at small non-zero lattice spacing (i.e. close to the classical continuum limit). Our approach is similar to the recent analytic work of O. Bär and I. Campos on the lattice version of Witten's global anomaly [9]. When combined with the preceding observation (1.3), our result implies that (1.4) is a necessary condition for gauge-invariance of the lattice chiral determinant in the overlap formulation, at least in the case where $\text{Map}(T^5, SU(N))$ contains maps with non-vanishing degree.⁷

It should be emphasised that global obstructions, and hence the results of this paper, are independent of the choice of phase in the overlap chiral determinant. In contrast, the consistent gauge anomaly for the overlap chiral determinant does depend on the phase choice. The consistent anomaly in the overlap formulation, and its classical continuum limit, has been previously studied in a number of works [10, 11, 12], although these have all involved some form of approximation (e.g. linearisation of the overlap) and/or assumptions (e.g. weak field, slowly varying field). No such approximations or assumptions are made in this paper.

The key question which these results lead on to (but which we do not pursue in this paper) is whether (1.4) is a *sufficient* condition for existence of a local anomaly-

⁷In higher dimensions there are cases where the maps all have vanishing degree yet the anomaly coefficient $d^{a_1 \cdots a_n}$ (the symmetrised trace of $T^{a_1}, \dots, T^{a_n}$) is non-vanishing. E.g. in dimension $2n = 6$ with gauge group $SU(3)$ one has $\pi_7 SU(3) = 0$ and $d^{a_1 a_2 a_3 a_4} \neq 0$, cf. p.472 of [3].

free lattice chiral gauge theory at non-zero lattice spacing. This is currently a topic of major interest and activity [5, 13, 14, 12, 15]. To proceed with this in practice, a specific phase choice must be made to begin with. (The standard choice is the so-called Brillouin–Wigner phase [2].) One can then try to “improve” the starting phase choice in various ways to get a local anomaly-free overlap when the cancellation condition holds. One practical approach is to average along the gauge orbits (i.e. the FNN mechanism); see [5] and the ref.’s therein.⁸ Another, more theoretical, approach is to reduce the problem of going from an arbitrary starting phase choice to one for which the (local) anomalies vanish to that of solving a system of finite-difference equations on the lattice [13, 14].⁹ In fact the integrability of these equations has been proved in the abelian case [13].¹⁰ There are strong indications that the same can be done in the nonabelian case [14, 12], although a complete proof of this has not yet been given. Quite recently, a practically-oriented analytic prescription for constructing anomaly-free non-compact chiral $U(1)$ gauge theory on the lattice has been given, starting from a adiabatic phase choice for the overlap [15].

In §2 we review the overlap construction of the chiral determinant. In §3 the lattice version of the global obstruction of Alvarez-Gaumé and Ginsparg is described, and is shown to reduce to the continuum obstruction in the classical continuum limit. The derivation of a key formula used to establish this is given separately in §4. This formula ((3.18) below) is due to Lüscher [14], and our detailed explicit derivation in §4 is intended to complement the rather brief argument in [14]. In §5 we make some concluding remarks. Some details of our calculations are given in an appendix.

⁸ The viability of this approach has been a topic of debate in the literature [16, 17, 18], although there is a body of evidence which is supportive of it —see, e.g., [19, 20].

⁹ The formulation of ref.’s [13, 14] (a functional integral formulation based on a lattice Dirac operator satisfying the Ginsparg-Wilson relation [21], which had been rediscovered outside of the overlap setting in the work of Hasenfratz and collaborators [22, 23]) is structurally identical to the overlap formulation after identifying the chiral fermion measures in the functional integral with the many-body groundstates in the overlap. More on this in §2, where the many-body groundstates are the “unit volume elements” in our terminology.

¹⁰ The argument in [13] relied on a result on the structure of the abelian axial anomaly [24], which has been further elucidated in [25].

2 Overlap construction of the chiral determinant on the lattice

The spacetime is taken to be the Euclidean 4-torus T^4 with fixed edge length L . (Again, the choice of dimension 4 is for concreteness and simplicity; the arguments and results in the following generalise straightforwardly to the T^{2n} case for arbitrary n .) We consider hyper-cubic lattices on T^4 with $2N$ sites along each edge and lattice spacing $a = L/2N$.¹¹ Given such a lattice, the space of lattice spinor fields $\psi(x)$ is denoted $\mathcal{C}$, and the space of lattice gauge fields $U_\mu(x)$ is denoted $\mathcal{U}$. The space $\mathcal{C}$ is finite-dimensional and comes equipped with an inner product:

$$\langle \psi_1, \psi_2 \rangle = a^4 \sum_x \psi_1(x)^* \psi_2(x) \quad (2.1)$$

With suitable boundary conditions, the covariant forward and backward finite difference operators $\frac{1}{a} \nabla_\mu^\pm$ act on $\mathcal{C}$ by

$$\nabla_\mu^+ \psi(x) = U_\mu(x) \psi(x + ae_\mu) - \psi(x) \quad (2.2)$$

$$\nabla_\mu^- \psi(x) = \psi(x) - U_\mu(x - ae_\mu)^{-1} \psi(x - ae_\mu) \quad (2.3)$$

e_μ denotes the unit vector in the positive μ -direction. We restrict to the case where $U_\mu(x)$ and $\psi(x)$ are periodic. This is the relevant case for considering the classical continuum limit with topologically trivial gauge fields. Since the chiral determinant vanishes in the topologically non-trivial case, this suffices for our purposes. Set $\nabla_\mu = \frac{1}{2}(\nabla_\mu^+ + \nabla_\mu^-)$; this operator is anti-hermitian with respect to the inner product (2.1) since $(\nabla_\mu^\pm)^* = -\nabla_\mu^\mp$. The Wilson–Dirac operator is now given by

$$D_{Wilson} = \frac{1}{a} \not{\nabla} + \frac{r}{2} a \left(\frac{1}{a^2} \Delta \right) \quad (2.4)$$

where $\not{\nabla} = \sum_\mu \gamma^\mu \nabla_\mu$ (the γ^μ 's are taken to be hermitian so $\not{\nabla}$ is anti-hermitian), $\Delta = \sum_\mu \nabla_\mu^- + \nabla_\mu^+ = \sum_\mu (\nabla_\mu^+)^* \nabla_\mu^+ = \sum_\mu (\nabla_\mu^-)^* \nabla_\mu^-$ (hermitian, positive) and $r > 0$ is the Wilson parameter. The hermitian operator

$$H(m) = \gamma_5 (a D_{Wilson} - rm) = \gamma_5 (\not{\nabla} + r(\frac{1}{2} \Delta - m)) \quad (2.5)$$

¹¹This N is of course not related to the N in $SU(N)$.

determines an orthogonal decomposition

$$\mathcal{C} = \mathcal{C}_+^{(m,U)} \oplus \mathcal{C}_-^{(m,U)} \quad (2.6)$$

where $\mathcal{C}_+^{(m,U)}$ and $\mathcal{C}_-^{(m,U)}$ are the subspaces spanned by the eigenvectors of $H(m)$ with positive and negative eigenvalues respectively. (We are restricting to the m, U for which $H(m)$ has no zero-modes.) These subspaces are characterised by $\epsilon(m) = \pm 1$ on $\mathcal{C}_\pm^{(m,U)}$ where

$$\epsilon(m) = \frac{H(m)}{\sqrt{H(m)^2}} \quad (2.7)$$

(the dependence on U has been suppressed). Noting that

$$\epsilon(m) = \frac{\frac{1}{|rm|}H(0) - \frac{m}{|m|}\gamma_5}{\sqrt{(\frac{1}{|rm|}H(0) - \frac{m}{|m|}\gamma_5)^2}} \rightarrow \gamma_5 \quad \text{for } m \rightarrow -\infty \quad (2.8)$$

we see that in the $m \rightarrow -\infty$ limit (2.6) reduces to the usual chiral decomposition

$$\mathcal{C} = \mathcal{C}_+ \oplus \mathcal{C}_- \quad (2.9)$$

independent of U . Set $m = 1$ (the canonical value; $0 < m < 2$ would suffice) and let $v_\pm$ and $w_\pm(U)$ be unit volume elements¹² on $\mathcal{C}_\pm$ and $\mathcal{C}_\pm^{(1,U)}$ respectively; these are unique up to phase factors. Then the lattice versions of the right- and left-handed chiral determinants in the overlap construction are, respectively,

$$\langle v_+, w_+(U) \rangle \quad (\text{right-handed}) \quad (2.10)$$

$$\langle v_-, w_-(U) \rangle \quad (\text{left-handed}) \quad (2.11)$$

(see [2] for background and motivation). The $w_\pm(U)$ are required to depend smoothly on U ; then the overlaps (2.10)–(2.11) are smooth in U . Note that a condition for non-vanishing overlaps is $\dim \mathcal{C}_\pm^{(1,U)} = \dim \mathcal{C}_\pm \equiv d$. The overlaps are unique up

¹²A vectorspace V determines vectorspaces $\Lambda^p V$ ($p=1, \dots, \dim V$): the exterior algebra (=Grassmann algebra) of V of degree p . An inner product in V induces an inner product in each $\Lambda^p V$. A “unit volume element on V ” is an element $v \in \Lambda^d V$ ($d = \dim V$) with $|v| = 1$. E.g. if $v_1, \dots, v_d$ is an orthonormal basis for V then $v_1 \wedge \dots \wedge v_d$ is a unit volume element. Since $\Lambda^d V$ is 1-dimensional, a unit volume element is unique up to $\pm$ if V is real, or up to a phase if V is complex.

to a phase factor, and their norms are gauge-invariant (an easy consequence of the gauge-covariance of $H(m)$, $\epsilon(m)$).

Remark 2.1. The construction of the overlaps (2.10)–(2.11) requires that $H(1)$ has no zero-modes. This can be guaranteed by imposing the condition [26, 27]

$$||1 - U(p)|| \leq 0.04 \quad \forall p \quad (2.12)$$

on the lattice gauge field U , where $U(p)$ is the product of the link variables around a plaquette p . This condition is automatically satisfied in the classical continuum limit since $1 - U(p_{x;\mu,\nu}) = -a^2 F_{\mu\nu}(x) + O(a^3)$.

We henceforth restrict $\mathcal{U}$ to be the space of lattice gauge fields satisfying (2.12).

Remark 2.2. The overlaps (2.10)–(2.11) are determined (up to a phase) solely by $\epsilon = \epsilon(1)$. The construction could be carried through given any hermitian operator ϵ with the property $\epsilon^2 = 1$. The norms of the resulting overlaps would be gauge-invariant provided ϵ is gauge-covariant.

Remark 2.3. The overlaps (2.10)–(2.11) can be written as¹³

$$\left(\frac{2}{a}\right)^d \langle v_+, w_+ \rangle = \langle v_+, \widehat{D} w_+ \rangle \equiv \det D_L \quad (2.13)$$

$$\left(\frac{2}{a}\right)^d \langle v_-, w_- \rangle = \langle v_-, \widehat{D} w_- \rangle \equiv \det D_R \quad (2.14)$$

where

$$D = \frac{1}{a} (1 + \gamma_5 \epsilon) \quad (2.15)$$

and $\epsilon = \epsilon(1)$ is given by (2.7). This follows easily from the facts that $(1 + \gamma_5 \epsilon)w = (1 \pm \gamma_5)w$ for $w \in \mathcal{C}_\pm^{(1,U)}$ and $(1 \pm \gamma_5)v = 2v$ for $v \in \mathcal{C}_\pm$. The relations (2.13)–(2.14) show how the overlaps can be viewed as chiral determinants in an analogous way to the continuum setting: Set $\hat{\gamma}_5 = \gamma_5(1 - aD) = -\epsilon$, then $\hat{\gamma}_5^2 = 1$ and $D\hat{\gamma}_5 = -\gamma_5 D$, which implies that D maps $\widehat{\mathcal{C}}_\mp := \mathcal{C}_\pm^{(1,U)}$ to $\mathcal{C}_\pm$. Thus, modulo the factors $(\frac{2}{a})^d$, the right-handed overlap can be viewed as a left-handed chiral determinant, and vice-versa, as

¹³ We are using the fact that a linear operator $D : W \rightarrow V$ induces linear operators $\widehat{D} : \Lambda^p W \rightarrow \Lambda^p V$ for all p , defined by $\widehat{D}(w_1 \wedge \cdots \wedge w_p) = Dw_1 \wedge \cdots \wedge Dw_p$. Note that if $W = V$ and $d = \dim V$ then $\widehat{D}(w_1 \wedge \cdots \wedge w_d) = \det D \cdot w_1 \wedge \cdots \wedge w_d$.

indicated in (2.13)–(2.14).¹⁴ In this form the overlap arises as the chiral determinant in Lüscher’s formulation [13, 14] after identifying the unit volume elements $v_{\pm}$ and $w_{\pm}$ (the many-body groundstates in the overlap) with the chiral fermion measures. These observations have been pointed out previously in [28, 29]. As mentioned there, it is easy to see that an operator D is of the form (2.15), with ϵ hermitian and $\epsilon^2 = 1$, if and only if it satisfies the the following two conditions:

$$D\gamma_5 + \gamma_5 D = aD\gamma_5 D \quad (\text{Ginsparg–Wilson relation [21]}) \quad (2.16)$$

$$D^* = \gamma_5 D \gamma_5 \quad (\gamma_5\text{–hermiticity}) \quad (2.17)$$

Also, clearly D is gauge-covariant if and only if ϵ is gauge-covariant. The operator (2.15), with $\epsilon = \epsilon(1)$ given by (2.7), is the Overlap Dirac operator introduced by Neuberger in [30]. It should also be mentioned that the Ginsparg–Wilson relation was rediscovered outside of the overlap setting in the work of P. Hasenfratz and collaborators—they considered a different solution, the so-called perfect Dirac operator [22, 23].

The nullspace of D is invariant under γ_5 (this follows from the GW relation (2.16): $D\psi=0 \Rightarrow D(\gamma_5\psi)=(aD\gamma_5D - \gamma_5D)\psi=0$) so $\text{index } D \equiv \text{Tr}(\gamma_5|_{\ker D})$ is well-defined, as was first noted in [23]. We only need to consider the lattice gauge fields U for which $\dim \mathcal{C}_{\pm}^{(1,U)} = \dim \mathcal{C}_{\pm} \equiv d$, since the overlaps vanish otherwise. As noted in [14], this corresponds to having $\text{index } D = 0$. Therefore, we henceforth take $\mathcal{U}$ to be the space of lattice gauge fields satisfying (2.12) and $\text{index } D = 0$.

¹⁴ A careful consideration of the overlap prescription shows that the overlaps $\langle v_{\pm}, w_{\pm} \rangle$ really should be multiplied by a factor $(\frac{2}{a})^d$ as in (2.13)–(2.14). These factors are physically irrelevant though: they appear both in the numerator and denominator in expressions for physical expectation values, and hence cancel out, and they do not affect anomalies since these only have to do with the phase of the overlaps. Nevertheless, they are relevant if one considers the chiral determinant on its own and wishes to use the lattice regularisation as an alternative to, e.g., zeta-regularisation.

3 Global obstructions to gauge-invariance of the Overlap

From now on we consider only the right-handed overlap $\langle v_+, w_+(U) \rangle$ (the situation for the left-handed overlap is analogous). A lattice version of the obstructions considered by Alvarez-Gaumé and Ginsparg is as follows. Let ϕ_θ be a family of lattice gauge transformations parameterised by $\theta \in S^1$. We can assume that the fermion content is specified by the fundamental representation of $SU(N)$; it will be clear from what follows that the general case is related to this case in the same way as in the continuum setting discussed in the introduction. If $U \in \mathcal{U}$ is a lattice gauge field for which the overlap $\langle v_+, w_+(U) \rangle$ is non-vanishing¹⁵ then the action of ϕ_θ on U determines a map

$$S^1 \rightarrow \mathbf{C} - \{0\} \quad , \quad \theta \mapsto \langle v_+, w_+(\phi_\theta \cdot U) \rangle \quad (3.1)$$

Since $|\langle v_+, w_+(U) \rangle|$ is gauge-invariant, we have $\langle v_+, w_+(\phi_\theta \cdot U) \rangle = e^{i\alpha(\theta)} \langle v_+, w_+(U) \rangle$ for some phase $\alpha(\theta)$, and the map (3.1) has integer winding number $W(\Phi, U) = \frac{1}{2\pi}(\alpha(1) - \alpha(0))$. Obviously, if the winding number is non-vanishing $\langle v_+, w_+(U) \rangle$ cannot be gauge-invariant. To see that this is a genuine obstruction to gauge-invariance we note that it is independent of the choice of $w_+(U)$: If $\tilde{w}_+(U)$ is another unit volume element on $\mathcal{C}_+^{(1,U)}$, smoothly varying with U , then $\tilde{w}_+(U) = e^{i\beta(U)} w_+(U)$ where the phase factor $e^{i\beta(U)}$ is smooth in U , and we have

$$\langle v_+, \tilde{w}_+(\phi_\theta \cdot U) \rangle = e^{i\alpha(\theta) + i\beta(\phi_\theta \cdot U)} \langle v_+, w_+(U) \rangle$$

Assuming that $\{\phi_\theta \cdot U\}_{\theta \in S^1}$ is a contractible circle in $\mathcal{U}$ (which is certainly true close to the classical continuum limit), it follows that this has the same winding number as (3.1) since $e^{i\beta(U)}$ is a smooth, non-vanishing, globally defined function of U . Hence the winding number $W(\Phi, U)$ is an obstruction to gauge-invariance of the overlap, independent of the choice of w_+ .

Our main result is that this obstruction coincides with the continuum one at small non-zero lattice spacing close to the classical continuum limit:

¹⁵ It can be seen from (2.13) that $\langle v_+, w_+(U) \rangle$ vanishes at the U for which D has zero-modes. Generically, these are isolated points in $\mathcal{U}$.

Theorem. If ϕ_θ is the restriction to the lattice of a family of continuum gauge transformations (also denoted ϕ_θ) and U is the lattice transcript of a topologically trivial continuum gauge field, then there is an $a_0 > 0$ (depending on the ϕ_θ 's and U) such that

$$W(\Phi, U) = \deg(\Phi) \quad \text{for all } a < a_0 \quad (3.2)$$

where $\deg(\Phi)$ is the degree of the continuum map $\Phi : T^5 \rightarrow SU(N)$ given by $\Phi(\theta, x) = \phi_\theta(x)$.

In light of the discussion in the introduction we conclude from this that, in the general case where the fermion content is specified by a general representation R of $SU(N)$, a necessary condition for existence of a gauge-invariant construction of the overlap is

$$d_R^{abc} = 0 \quad (3.3)$$

The remainder of the paper is concerned with the proof of the above theorem. We start by expressing the obstruction as

$$\begin{aligned} W(\Phi, U) &= \frac{1}{2\pi i} \int_0^1 d\theta \frac{d}{d\theta} \log \langle v_+, w_+(\phi_\theta \cdot U) \rangle \\ &= \frac{1}{2\pi i} \int_{\mathcal{S}^1} d \log \langle v_+, w_+ \rangle \end{aligned} \quad (3.4)$$

where $\mathcal{S}^1$ denotes the circle $\{\phi_\theta \cdot U\}_{\theta \in \mathcal{S}^1}$ in $\mathcal{U}$ and d denotes the exterior derivative on functions (or more generally, differential forms) on $\mathcal{U}$. After noting that

$$dw_+ = \langle w_+, dw_+ \rangle w_+ + (dw_+)_{\perp} \quad (3.5)$$

where $(dw_+)_{\perp}$ denotes the projection of dw_+ onto the orthogonal complement of w_+ in $\Lambda^d \mathcal{C}$, one finds [5]

$$d \log \langle v_+, w_+ \rangle = \frac{\langle v_+, (dw_+)_{\perp} \rangle}{\langle v_+, w_+ \rangle} + \langle w_+, dw_+ \rangle \quad (3.6)$$

The first term on the right-hand side of (3.6) can be re-expressed as

$$\frac{\langle v_+, (dw_+)_{\perp} \rangle}{\langle v_+, w_+ \rangle} = \text{Tr}(dDD^{-1}P_+) \quad (3.7)$$

where $P_+ = \frac{1}{2}(1 + \gamma_5)$ is the projection onto $\mathcal{C}_+$. This is a straightforward consequence of (2.13) and relations noted in [13, 14]; for completeness we provide a derivation in the appendix. Then

$$d \log \langle w_+, w_+ \rangle = \text{Tr}(dDD^{-1}P_+) + \langle w_+, dw_+ \rangle. \quad (3.8)$$

Set $w_+^\theta = w_+(\phi_\theta \cdot U)$ and let D_θ denote D with lattice gauge field $\phi_\theta \cdot U$. The gauge-covariance of D gives $\frac{d}{d\theta}D_\theta = [\frac{d\phi_\theta}{d\theta}\phi_\theta^{-1}, D_\theta]$, leading to [13, 14]

$$\begin{aligned} \text{Tr}(\frac{d}{d\theta}D_\theta D_\theta^{-1}P_+) &= \text{Tr}\left(\frac{d\phi_\theta}{d\theta}\phi_\theta^{-1}P_+\right) - \text{Tr}\left(D_\theta \frac{d\phi_\theta}{d\theta}\phi_\theta^{-1}D_\theta^{-1}P_+\right) \\ &= \text{Tr}\left(\frac{d\phi_\theta}{d\theta}\phi_\theta^{-1}(P_+ - \hat{P}_-)\right) \\ &= -\frac{1}{2}a\text{Tr}\left(\frac{d\phi_\theta}{d\theta}\phi_\theta^{-1}\gamma_5 D_\theta\right) = -\frac{1}{2}a\text{Tr}\left(\phi_\theta^{-1}\frac{d\phi_\theta}{d\theta}\gamma_5 D_1\right) \end{aligned} \quad (3.9)$$

where we have used the fact that $P_+D = D\hat{P}_-$ where $\hat{P}_- = \frac{1}{2}(1 - \hat{\gamma}_5)$, $\hat{\gamma}_5 = \gamma_5(1 - aD)$.

Substituting (3.8) into (3.4) and using (3.9) we get

$$2\pi i W(\Phi, U) = \int_{S^1} \text{Tr}(dDD^{-1}P_+) + \langle w_+, dw_+ \rangle \quad (3.10)$$

$$= \int_0^1 d\theta \left(-\frac{1}{2}a\text{Tr}\left(\phi_\theta^{-1}\frac{d\phi_\theta}{d\theta}\gamma_5 D_1\right) + \langle w_+^\theta, \frac{dw_+^\theta}{d\theta} \rangle \right) \quad (3.11)$$

We have derived this relation under the assumption that the overlap is non-vanishing for U , or equivalently, that the Overlap Dirac operator D with lattice gauge field U has no zero-modes. By construction $W(\Phi, U)$ is clearly smooth, and therefore locally constant, in such U . But it is ill-defined at the (generically isolated) points in $\mathcal{U}$ where the overlap vanishes. One such point is the trivial field $U = 1$ (in this case the zero-momentum spinors with definite chirality are zero-modes for D). However, the right-hand side of (3.11) is clearly smooth in U for all $U \in \mathcal{U}$ (since D is smooth in the lattice gauge field when (2.12) is satisfied [26]), and must therefore be a locally constant function of U for all $U \in \mathcal{U}$. In the continuum, any topologically trivial gauge field can be continuously deformed to the trivial field. It follows that when the lattice spacing a is sufficiently small, the lattice transcript U can be continuously deformed to the trivial lattice gauge field (using the lattice transcript of the continuum path). Therefore, to prove the theorem it suffices to show that there is an $a_0 > 0$ such that

$$W(\Phi) = \deg(\Phi) \quad \text{for all } a < a_0 \quad (3.12)$$

where $W(\Phi)$ denotes the right-hand side of (3.11) with trivial field $U = 1$. In this case D_1 acts trivially in colour space, and since $\phi_\theta^{-1} \frac{d\phi_\theta}{d\theta}$ is in the Lie algebra of $SU(N)$ the trace over colour indices in the first term in (3.11) vanishes, resulting in

$$\begin{aligned} 2\pi i W(\Phi) &= \int_0^1 d\theta \langle w_+^\theta, \frac{d}{d\theta} w_+^\theta \rangle \\ &= \int_{\mathcal{S}^1} \langle w_+, dw_+ \rangle \end{aligned} \quad (3.13)$$

A calculation gives

$$\begin{aligned} d\langle w_+, dw_+ \rangle &= \langle dw_+, dw_+ \rangle \\ &= \text{Tr}(PdPdP) \end{aligned} \quad (3.14)$$

where

$$P \equiv \hat{P}_- = \frac{1}{2}(1 - \hat{\gamma}_5) = \frac{1}{2}(1 + \epsilon) \quad (3.15)$$

with $\epsilon = \epsilon(1)$ given by (2.7). The last equality in (3.14) is derived in the appendix. A simpler version of it (originating in [31]) was used in [5]; the same relation in a different guise was subsequently noted in [13, 14], and in more detail in [11]. Now, by Stokes theorem, if $\mathcal{B}^2$ is a disc in $\mathcal{U}$ with boundary $\mathcal{S}^1$ it follows that

$$\begin{aligned} 2\pi i W(\Phi) &= \int_{\mathcal{B}^2} \text{Tr}(PdPdP) \\ &= \int_{\mathcal{B}^2} \text{Tr}(P[\partial_\theta P, \partial_t P]) d\theta dt \end{aligned} \quad (3.16)$$

where (θ, t) are taken to be polar coordinates on a unit disc B^2 parameterising $\mathcal{B}^2$. We take $\mathcal{B}^2$ to be the lattice transcript of a disc-family of continuum gauge fields, also denoted $\mathcal{B}^2$, given by

$$A^{(\theta, t)} = f(t) \phi_\theta d_x \phi_\theta^{-1} \quad (\theta, t) \in B^2 \quad (3.17)$$

where $f(t)$ is an arbitrary smooth function equal to 1 in a neighbourhood of $t=1$ and vanishing in a neighbourhood of $t=0$. The lattice transcript $U^{(\theta, t)}$ has the property $U^{(\theta, 1)} = \phi_\theta \cdot 1$ so the boundary of $\mathcal{B}^2$ is $\mathcal{S}^1$ as required in (3.16). Note that (3.16) is manifestly independent of the choice of $w_+(U)$, i.e. independent of the choice of phase in the overlap.

A general formula for the classical continuum limit of the integrand in (3.16) has been given by Lüscher in [14]: If $U^{(s,t)}$ is the lattice transcript of a family $A^{(s,t)}$ of continuum gauge fields, and $P = P^{(s,t)}$ is the corresponding family of projection operators (given by (3.15)), then

$$\lim_{a \rightarrow 0} \text{Tr}(P[\partial_s P, \partial_t P]) = \frac{-1}{32\pi^2} \int_{T^4} d^4x \epsilon_{\mu\nu\rho\sigma} d^{abc} \partial_s A_\mu^a(x) \partial_t A_\nu^b(x) F_{\rho\sigma}^c(x). \quad (3.18)$$

Using this, (3.12) follows easily from (3.16) as we will see below, thereby proving the theorem. In [14] the locality, smoothness and symmetry properties of P were used to show that the limit in (3.18) exists and is given by the integral over T^4 of a polynomial in the gauge fields and its derivatives. However, the explicit form of this polynomial (i.e. the integrand on the right-hand side of (3.18)) was not obvious, at least to the present author, from the brief argument in [14]. Since (3.18) is an important formula in this context (it was also a key ingredient in the arguments of ref.'s [14] and [9]) we will give a detailed, explicit derivation of it in §4. This is intended to complement the brief argument in [14].

By (3.18), the classical continuum limit of (3.16) is

$$\lim_{a \rightarrow 0} 2\pi i W(\Phi) = \frac{-1}{32\pi^2} \int_{B^2 \times T^4} d\theta dt d^4x \epsilon_{\mu\nu\rho\sigma} d^{abc} \partial_\theta A_\mu^a(x) \partial_t A_\nu^b(x) F_{\rho\sigma}^c(x). \quad (3.19)$$

with $A = A^{(\theta,t)}$ given by (3.17). It remains to show that this is equal to $2\pi i \deg(\Phi)$. Then, since $W(\Phi)$ is integer, we can conclude that $W(\Phi) = \deg(\Phi)$ for all lattice spacings a smaller than some $a_0 > 0$ and the theorem is proved. The right-hand side of (3.18) can be shown to equal $2\pi i \deg(\Phi)$ by a direct calculation [9], but it is easier and perhaps more illuminating to proceed indirectly as follows. We view the family $A^{(\theta,t)}$ as a gauge field on $B^2 \times T^4$:

$$\mathcal{A}(x, \theta, t) = A_\mu^{(\theta,t)}(x) dx^\mu = f(t) \phi_\theta(x) \partial_\mu \phi_\theta^{-1}(x) dx^\mu \quad (3.20)$$

and define another gauge field on $\tilde{B}^2 \times T^4$ by

$$\tilde{\mathcal{A}}(x, \theta, s) = -f(s) \partial_\theta \phi_\theta^{-1}(x) \phi_\theta(x) d\theta \quad (3.21)$$

where $\tilde{B}^2$ is another copy of the unit disc, with polar coordinates (θ, s) . $B^2 \times T^4$ and $\tilde{B}^2 \times T^4$ can be glued together along their common boundary $S^1 \times T^4$ to get the

closed manifold $S^2 \times T^4$. On the common boundary $S^1 \times T^4$ the fields $\mathcal{A}$ and $\tilde{\mathcal{A}}$ are related by a gauge transformation:

$$\tilde{\mathcal{A}} = \Phi^{-1} \cdot \mathcal{A} = \Phi^{-1} \mathcal{A} \Phi + \Phi^{-1} (d_\theta + d_x) \Phi$$

where $\Phi : \mathcal{S}^1 \times T^4 \rightarrow SU(N)$ is given by $\Phi(\theta, x) = \phi_\theta(x)$. Therefore $\mathcal{A}$ and $\tilde{\mathcal{A}}$ constitute a gauge field $\hat{\mathcal{A}}$ on an $SU(N)$ bundle over $S^2 \times T^4$ with topological charge $-\deg(\Phi)$. The topological charge is also given by the integral of the Chern character over $S^2 \times T^4$, thus

$$\begin{aligned} -\deg(\Phi) &= \frac{i^3}{(2\pi)^3 3!} \int_{S^2 \times T^4} \text{tr} \hat{\mathcal{F}}^3 \\ &= \frac{i^3}{(2\pi)^3 3!} \left[\int_{B^2 \times T^4} \text{tr} \mathcal{F}^3 - \int_{\tilde{B}^2 \times T^4} \text{tr} \tilde{\mathcal{F}}^3 \right] \end{aligned} \quad (3.22)$$

The second term vanishes: $\tilde{\mathcal{F}}^3 = 0$ since $\tilde{\mathcal{A}}$ only involves the 1-form $d\theta$. Regarding the first term, from $\mathcal{F} = (d_x + d_\theta + d_t)\mathcal{A} + \mathcal{A} \wedge \mathcal{A}$ we get

$$\begin{aligned} \text{tr} \mathcal{F}^3 &= \text{tr}(d_\theta A d_t A F + d_t A d_\theta A F + d_\theta A F d_t A + d_t A F d_\theta A + F d_\theta A d_t A + F d_t A d_\theta A) \\ &= 3 \left(\frac{d^{abc}}{2} \right) d_\theta A^a d_t A^b F^c \end{aligned}$$

After substituting this for $\text{tr} \mathcal{F}^3$ in (3.22) we see from (3.19) that $\lim_{a \rightarrow 0} W(\Phi) = \deg(\Phi)$ as required.

4 A detailed derivation of Lüscher's formula

In this section we give a detailed, explicit derivation of the formula (3.18). We begin by noting that

$$\begin{aligned} \text{Tr}(P[\partial_s P, \partial_t P]) &= \frac{1}{8} \text{Tr}(\epsilon[\partial_s \epsilon, \partial_t \epsilon]) \\ &= \frac{1}{4} \text{Tr}(\epsilon \partial_s \epsilon \partial_t \epsilon) \end{aligned} \quad (4.1)$$

Setting $H = H(1)$ we have

$$\epsilon = \frac{H}{\sqrt{H^2}} = \frac{\gamma_5 X}{\sqrt{X^* X}} \quad , \quad X = a\gamma_5 H = \not{X} + r(\tfrac{1}{2}\Delta - 1) \quad (4.2)$$

and a calculation using (4.1) gives

$$\begin{aligned} & \text{Tr}(P[\partial_s P, \partial_t P]) \\ &= \frac{1}{4} \text{Tr} \left(\frac{\gamma_5 X}{\sqrt{X^* X}} \partial_s (\gamma_5 X) \frac{1}{\sqrt{X^* X}} \partial_t (\gamma_5 X) \frac{1}{\sqrt{X^* X}} \right) \end{aligned} \quad (4.3)$$

$$+ \frac{1}{4} \text{Tr} \left(\partial_s \left(\frac{1}{\sqrt{X^* X}} \right) \partial_t (\gamma_5 X) \right) + \frac{1}{4} \text{Tr} \left(\frac{1}{X^* X} \partial_s (\gamma_5 X) \partial_t \sqrt{X^* X} \right) \quad (4.4)$$

$$- \frac{1}{4} \text{Tr} \left(\partial_s \left(\frac{1}{\sqrt{X^* X}} \right) \sqrt{X^* X} \gamma_5 X \partial_t \left(\frac{1}{\sqrt{X^* X}} \right) \right) \quad (4.5)$$

We will see below that (4.4) $\sim O(a)$ and (4.5) $= \text{Symm} + O(a)$ where *Symm* is symmetric under interchange of ∂_s and ∂_t . Since $\text{Tr}(P[\partial_s P, \partial_t P])$ is antisymmetric under this interchange it follows that

$$\begin{aligned} \text{Tr}(P[\partial_s P, \partial_t P]) &= \frac{1}{8} \left[\text{Tr} \left(\frac{\gamma_5 X}{\sqrt{X^* X}} \partial_s (\gamma_5 X) \frac{1}{\sqrt{X^* X}} \partial_t (\gamma_5 X) \frac{1}{\sqrt{X^* X}} \right) \right. \\ &\quad \left. - \text{Tr} \left(\frac{\gamma_5 X}{\sqrt{X^* X}} \partial_t (\gamma_5 X) \frac{1}{\sqrt{X^* X}} \partial_s (\gamma_5 X) \frac{1}{\sqrt{X^* X}} \right) \right] + O(a) \end{aligned} \quad (4.6)$$

To prove these statements, and evaluate the $a \rightarrow 0$ limit of (4.6), we use the fact that the lattice transcript¹⁶

$$U_\mu(x) = T \exp \left(\int_0^1 a A_\mu(x + (1 - \tau) a e_\mu) d\tau \right) \quad (4.7)$$

can be expanded in powers of a as

$$U_\mu(x) = \sum_{n=0}^{\infty} a^n \int_{0 < \tau_1 < \dots < \tau_n < 1} d\tau_n \cdots d\tau_1 A_\mu(x, \tau_n) \cdots A_\mu(x, \tau_1) \quad (4.8)$$

where $A_\mu(x, \tau) = A_\mu(x + (1 - \tau) a e_\mu)$. Since $A_\mu(x)$ is a smooth function on the closed manifold T^4 there is a finite K such that $|A_\mu(x)| < K$ for all x, μ . Then the norm of the integral in the n 'th term of (4.8) is bounded by $\frac{1}{n!} K^n$, so (4.8) is norm-convergent for all a . The inverse $U_\mu(x)^{-1}$ also has an expansion in powers of a : using the fact that $U_\mu(x)^{-1}$ is the parallel transport from x to $x + a e_\mu$ specified by A we see that $U_\mu(x)^{-1}$ is given by the right-hand side of (4.8) with $A_\mu(x, \tau) = A_\mu(x + \tau a e_\mu)$.

Substituting these expansions in (2.2)–(2.3) gives an expansion $\nabla_\mu^\pm = \sum_{n=0}^{\infty} a^n (\nabla_\mu^\pm)_n$. This in turn gives an expansion $X = \sum_{n=0}^{\infty} a^n X_n$. It is not difficult to show that the

¹⁶ Here $A = A^{(s,t)}$ and $U = U^{(s,t)}$ depend smoothly on the two parameters (s, t) .

$||(\nabla_\mu^\pm)_n||$'s and the $||X_n||$'s have a finite bound K' independent of a and n , so the expansions are norm-convergent when a is sufficiently small.

To expand $1/\sqrt{X^*X}$ we note that

$$X^*X = L + V \quad (4.9)$$

where

$$L = -\nabla^2 + r^2(\frac{1}{2}\Delta - m)^2 \quad (4.10)$$

$$V = V^{(1)} + V^{(2)} \quad (4.11)$$

$$V^{(1)} = \frac{1}{2}r\gamma^\mu[\nabla_\mu, \sum_\nu \nabla_\nu^+ - \nabla_\nu^-] \quad , \quad V^{(2)} = -\frac{1}{4}[\gamma^\mu, \gamma^\nu][\nabla_\mu, \nabla_\nu] \quad (4.12)$$

Just as for X , the expansion (4.8) leads to expansions $X^*X = \sum_{n=0}^\infty a^n(X^*X)_n$, $L = \sum_{n=0}^\infty a^n L_n$, $V = \sum_{n=0}^\infty a^n V_n$ where the $||(X^*X)_n||$'s, $||L_n||$'s and $||V_n||$'s again have a finite bound independent of a and n . Furthermore, explicit calculations show that

$$[\nabla_\mu^+, \nabla_\nu^+]\psi(x) = (-a^2 F_{\mu\nu}(x) + O(a^3))\psi(x + ae_\mu + ae_\nu) \quad (4.13)$$

$$[\nabla_\mu^+, \nabla_\nu^-]\psi(x) = (-a^2 F_{\mu\nu}(x) + O(a^3))\psi(x + ae_\mu - ae_\nu) \quad (4.14)$$

$$[\nabla_\mu^-, \nabla_\nu^+]\psi(x) = (-a^2 F_{\mu\nu}(x) + O(a^3))\psi(x - ae_\mu + ae_\nu) \quad (4.15)$$

$$[\nabla_\mu^-, \nabla_\nu^-]\psi(x) = (-a^2 F_{\mu\nu}(x) + O(a^3))\psi(x - ae_\mu - ae_\nu) \quad (4.16)$$

It follows that $V_0 = V_1 = 0$, i.e. the expansion of V starts with the a^2 term, hence $||V|| \sim O(a^2)$. The leading term $a^2 V_2$ is explicitly given (mod $O(a^3)$) by substituting (4.13)–(4.16) into (4.11)–(4.12). We note from this that $V_2 = V_2^b T^b$ where the T^b 's are the generators of the Lie algebra of $SU(N)$ and the V_2^b 's are trivial in colour space.

From (4.8) we also get expansions of $\partial_s U_\mu(x)$ and $\partial_s U_\mu(x)^{-1} = U_\mu(x)^{-1} \partial_s U_\mu(x) U_\mu(x)^{-1}$ in powers of a , leading to an expansion $\partial_s X = \sum_{n=0}^\infty a^n (\partial_s X)_n$ and expansions of $\partial_s(X^*X)$, $\partial_s L$, and $\partial_s V$. Note that these begin with the order a term, i.e. $(\partial_s X)_0 = (\partial_s(X^*X))_0 = (\partial_s L)_0 = (\partial_s V)_0 = 0$. For later use we also note the following: (1) The lowest order term in the expansion of $\partial_s U_\mu(x)$ (or $\partial_s U_\mu(x)^{-1}$) is $a A_\mu^b(x) T^b$ (or $-a A_\mu^b(x) T^b$). (2) Applying ∂_s to (4.13)–(4.16) results in $F_{\mu\nu}(x) \rightarrow \partial_s F_{\mu\nu}(x)$, so

$(\partial_s V)_0 = (\partial_s V)_1 = 0$, $\|\partial_s V\| \sim O(a^2)$, and $(\partial_s V)_2 = (\partial_s V)_2^b T^b$ where the $(\partial_s V)_2^b$'s are trivial in colour space.

Note that the γ^μ 's in (4.9) are all contained in V . The hermitian positive operator L is trivial in Dirac indices and the lowest order term L_0 in its expansion is diagonal with respect to the plane wave basis $\{e^{iak \cdot x}\}$; the diagonal elements are

$$L_0(ak) = \sum_{\mu} \sin^2(ak_{\mu}) + r^2 \left(-1 + \sum_{\mu} 1 - \cos(ak_{\mu}) \right)^2 \quad (4.17)$$

From this we see that there is a $b > 0$ independent of a such that $L_0 > 2b$. Then, by taking a to be sufficiently small, we can achieve $L > b$ and $\|V\| < \frac{1}{2}b$, in which case $1/\sqrt{X^*X}$ can be expanded as follows:

$$\begin{aligned} \frac{1}{\sqrt{X^*X}} &= \int_{-\infty}^{\infty} \frac{d\sigma}{\pi} \frac{1}{X^*X + \sigma^2} \\ &= \int_{-\infty}^{\infty} \frac{d\sigma}{\pi} \left(\frac{1}{1 + (L + \sigma^2)^{-1}V} \right) \left(\frac{1}{L + \sigma^2} \right) \\ &= \int_{-\infty}^{\infty} \frac{d\sigma}{\pi} \sum_{k=0}^{\infty} (-1)^k ((L + \sigma^2)^{-1}V)^k \frac{1}{L + \sigma^2}. \end{aligned} \quad (4.18)$$

For all p we have

$$\left\| \sum_{k=0}^p (-1)^k ((L + \sigma^2)^{-1}V)^k \frac{1}{L + \sigma^2} \right\| < \sum_{k=0}^{\infty} \left(\frac{1}{b} \left(\frac{1}{2}b \right) \right)^k \frac{1}{b + \sigma^2} = \frac{2}{b + \sigma^2}$$

Since the integral of this over $(-\infty, \infty)$ is finite, the integral and sum in (4.18) can be interchanged, resulting in a norm-convergent expansion in powers of V :

$$(X^*X)^{-1/2} = \sum_{k=0}^{\infty} \int_{-\infty}^{\infty} \frac{d\sigma}{\pi} (-1)^k ((L + \sigma^2)^{-1}V)^k (L + \sigma^2)^{-1} \quad (4.19)$$

Since $\|V\| \sim O(a^2)$ the k 'th term in the sum is $\sim O(a^{2k})$ and we conclude that

$$(X^*X)^{-1/2} = \sum_{k=0}^p \int_{-\infty}^{\infty} \frac{d\sigma}{\pi} (-1)^k ((L + \sigma^2)^{-1}V)^k (L + \sigma^2)^{-1} + R_{p+1} \quad (4.20)$$

where $\frac{1}{a^{2p}} \|R_{p+1}\| \rightarrow 0$ for $a \rightarrow 0$. Similarly, we find

$$\partial_s (X^*X)^{-1/2} = \sum_{k=0}^p \int_{-\infty}^{\infty} \frac{d\sigma}{\pi} (-1)^k \partial_s \left(((L + \sigma^2)^{-1}V)^k (L + \sigma^2)^{-1} \right) + \partial_s R_{p+1} \quad (4.21)$$

where $\frac{1}{a^{2p}} \|\partial_s R_{p+1}\| \rightarrow 0$ for $a \rightarrow 0$.

The bound $\|\nabla_\mu^\pm\| \leq 2$ and triangle inequalities lead to an a -independent upper bound $L < b_1$. Using this, the operator $(L + \sigma^2)^{-1}$ in (4.20) can be expanded as

$$\begin{aligned} \frac{1}{L + \sigma^2} &= \left(\frac{1}{b_1 + \sigma^2} \right) \left(\frac{1}{1 - \frac{b_1 - L}{b_1 + \sigma^2}} \right) \\ &= (b_1 + \sigma^2)^{-1} \sum_{m=0}^{\infty} (b_1 + \sigma^2)^{-m} (b_1 - L)^m \end{aligned} \quad (4.22)$$

Substituting the expansion $L = \sum_{n=0}^{\infty} a^n L_n$ in (4.22), and then substituting in (4.20) the resulting expansion of $(L + \sigma^2)^{-1}$, along with the expansion $V = \sum_{n=2}^{\infty} a^n V_n$, we get an expansion $(X^* X)^{-1/2} = \sum_{n=0}^{\infty} a^n (X^* X)_n^{-1/2}$. Similarly, after applying ∂_s to (4.22) and substituting the resulting expansion in (4.21), we get an expansion $\partial_s (X^* X)^{-1/2} = \sum_{n=1}^{\infty} a^n (\partial_s (X^* X)^{-1/2})_n$ (note $(\partial_s (X^* X)^{-1/2})_0 = 0$). These, together with the expansions of X and ∂X , lead to expansions $\mathcal{O} = \sum_{n=0}^{\infty} a^n \mathcal{O}_n$ of the operators in (4.3)–(4.5). It can be shown that there is a finite $\widetilde{K}$ independent of a and n such that $\|\mathcal{O}_n\| < \widetilde{K}$ for these operators. This technical result will be presented elsewhere [32]¹⁷. This implies that the aforementioned operator expansions are all norm-convergent when a is sufficiently small. An immediate consequence is the following: In the resulting expressions $\sum_{n=0}^{\infty} a^n \text{Tr}(\mathcal{O}_n)$ for (4.3)–(4.5) the part $\sum_{n=5}^{\infty} a^n \text{Tr}(\mathcal{O}_n)$ vanishes in the $a \rightarrow 0$ limit. To see this, let $\{\psi_j\}_{j=1, \dots, \mathcal{N}}$ be an arbitrary orthonormal basis for $\mathcal{C}$; then

$$\begin{aligned} \left| \sum_{n \geq 5} a^n \text{Tr}(\mathcal{O}_n) \right| &\leq \sum_{n \geq 5, j} a^n |\langle \psi_j, \mathcal{O}_n \psi_j \rangle| \\ &\leq \sum_{n \geq 5, j} a^n \widetilde{K} = a^4 \mathcal{N} \widetilde{K} \sum_{n=0}^{\infty} a^n = 4L^4 \widetilde{K} \frac{a}{1-a} \rightarrow 0 \quad \text{for } a \rightarrow 0 \end{aligned} \quad (4.23)$$

where we have used the fact that $\mathcal{N} \equiv \dim \mathcal{C} = 4(L/a)^4$. This shows that we only need to consider the a^n terms with $n \leq 4$ in the expansions $\sum_{n=0}^{\infty} a^n \text{Tr}(\mathcal{O}_n)$ of (4.3)–(4.5).

To proceed with the determination of (4.3)–(4.5) in the $a \rightarrow 0$ limit, we note the following: (i) Due to the presence of an odd number of γ_5 's, terms in the expansions

¹⁷ It relies on the fact that $A_\mu(x)$ is periodic. The general (i.e. topologically non-trivial) case is more complicated.

involving a product of less than 4 γ^μ 's vanish. (ii) L , $V^{(1)}$ and $V^{(2)}$ are of order 0, 1 and 2 respectively in the γ^μ -matrices, c.f. (4.12). (iii) In the expansion $\mathcal{O} = \sum_{n=0}^{\infty} a^n \mathcal{O}_n$ of any operator $\mathcal{O}$ constructed from the $\nabla_\mu^\pm$'s, the term $\mathcal{O}_0$ is independent of the gauge field, so $(\partial_s \mathcal{O})_0 = 0$. Hence non-vanishing terms in such expansions are at least $O(a)$ for $a \rightarrow 0$. (iv) As we have seen in (4.23), terms in the operator expansions which are of order ≥ 5 in a give vanishing contributions in the $a \rightarrow 0$ limit.

At this point we can derive the postulated formula (4.6). Consider the first trace in (4.4): After substituting the expansion (4.19) for $(X^* X)^{-1/2}$ only the terms with at least two V 's are non-vanishing after taking the trace over spinor indices. Since $\|V\| \sim O(a^2)$, such terms are all of order ≥ 4 in a . Since $\partial_t(\gamma_5 X) \sim O(a)$ it follows that the terms in the expansion of $\partial_s(X^* X)^{-1/2} \partial_t(\gamma_5 X)$ which are non-vanishing after taking the trace are all of order ≥ 5 in the $a \rightarrow 0$ limit, so (4.4) $\sim O(a)$ as claimed.

We now consider the trace (4.5). First, note from (4.20)–(4.22) that the lowest order term involving V (or containing γ^μ 's) in the expansion of $(X^* X)^{-1/2}$ is

$$- \int_{-\infty}^{\infty} \frac{d\sigma}{\pi} (L_0 + \sigma^2)^{-1} a^2 V_2 (L_0 + \sigma^2)^{-1}. \quad (4.24)$$

Using (4.22) and the fact that $[L_0, V] \sim O(a)$ we find that, modulo an $O(a)$ term, this is

$$- a^2 \int_{-\infty}^{\infty} \frac{d\sigma}{\pi} (L_0 + \sigma^2)^{-2} V_2 = - a^2 L_0^{-3/2} V_2 \int_{-\infty}^{\infty} \frac{d\sigma}{\pi} \frac{1}{(1 + \sigma^2)^2} = - \frac{1}{2} a^2 L_0^{-3/2} V_2 \quad (4.25)$$

Similarly, the lowest order term containing γ^μ 's in the expansion $\partial_s(X^* X)^{-1/2}$ is $-\frac{1}{2} a^2 L_0^{-3/2} (\partial_s V)_2$. To simplify the notation in the following, we write $\partial_s \mathcal{O}_n$ for $(\partial_s \mathcal{O})_n$. Now, using (i)–(iv) above, we find

$$\begin{aligned} & \text{Tr} \left(\partial_s (X^* X)^{-1/2} (X^* X)^{1/2} \gamma_5 X \partial_t (X^* X)^{-1/2} \right) \\ &= a^4 \text{Tr} \left(\partial_s (X^* X)_2^{-1/2} (X^* X)_0^{1/2} \gamma_5 X_0 \partial_t (X^* X)_2^{-1/2} \right) + O(a) \\ &= \frac{a^4}{4} \text{Tr} \left(L_0^{-3/2} \partial_s V_2 L_0^{1/2} \gamma_5 X_0 L_0^{-3/2} \partial_t V_2 \right) + O(a) \end{aligned} \quad (4.26)$$

where we have also used $(X^* X)_0 = L_0$. We now supplement (i)–(iv) above with the following observation (v): V_2 and $(\partial V)_2$ commute with L_0 modulo an $O(a)$ term,

and commute with $\gamma_5 X_0$ modulo an $O(a)$ term and a term of order 1 in the γ^μ 's. It follows that, modulo an $O(a)$ term, (4.26) is symmetric under interchange of ∂_s and ∂_t as claimed. This proves the previously stated symmetry property of (4.5), thereby completing the derivation of (4.6).

Turning now to the traces in (4.6), similar arguments to the preceding give

$$\begin{aligned}
& \text{Tr} \left(\gamma_5 X (X^* X)^{-1/2} \partial_s (\gamma_5 X) (X^* X)^{-1/2} \partial_t (\gamma_5 X) (X^* X)^{-1/2} \right) \\
&= a^4 \text{Tr} \left(\gamma_5 X_0 (X^* X)_2^{-1/2} \partial_s (\gamma_5 X)_1 (X^* X)_0^{-1/2} \partial_t (\gamma_5 X)_1 (X^* X)_0^{-1/2} \right) \\
&+ a^4 \text{Tr} \left(\gamma_5 X_0 (X^* X)_0^{-1/2} \partial_s (\gamma_5 X)_1 (X^* X)_2^{-1/2} \partial_t (\gamma_5 X)_1 (X^* X)_0^{-1/2} \right) \\
&+ a^4 \text{Tr} \left(\gamma_5 X_0 (X^* X)_0^{-1/2} \partial_s (\gamma_5 X)_1 (X^* X)_0^{-1/2} \partial_t (\gamma_5 X)_1 (X^* X)_2^{-1/2} \right) + O(a) \\
&= -\frac{1}{2} a^4 (t^{cab} + t^{acb} + t^{abc}) \text{Tr} \left(L_0^{-5/2} \gamma_5 X_0 \partial_s (\gamma_5 X)_1^a \partial_t (\gamma_5 X)_1^b V_2^c \right) + O(a) \quad (4.27)
\end{aligned}$$

where we have set $t^{abc} = \text{tr}(T^a T^b T^c)$. Modulo an $O(a)$ term, (4.27) is antisymmetric under interchange of $\partial_s (\gamma_5 X)_1^a$ and $\partial_t (\gamma_5 X)_1^b$. To see this, note that $\partial_s (X^* X)_1^a = \gamma_5 X_0 \partial_s (\gamma_5 X)_1^a + \partial_s (\gamma_5 X)_1^a \gamma_5 X_0$. Since $(X^* X)_1 = L_1$ does not contain γ^μ 's, it follows that $\gamma_5 X_0 \partial_s (\gamma_5 X)_1^a$ can be replaced by $-\partial_s (\gamma_5 X)_1^a \gamma_5 X_0$ in (4.27). The claimed antisymmetry then follows from the cyclicity of the trace after using (v) above. Taking this into account in (4.6), and noting $\frac{1}{2} d^{abc} = t^{cab} + t^{cba} = t^{acb} + t^{bca} = t^{abc} + t^{bac}$, we get

$$\text{Tr}(P[\partial_s P, \partial_t P]) = -\frac{3}{32} a^4 d^{abc} \text{Tr} \left(L_0^{-5/2} \gamma_5 X_0 \partial_s (\gamma_5 X)_1^a \partial_t (\gamma_5 X)_1^b V_2^c \right) + O(a) \quad (4.28)$$

We calculate

$$\begin{aligned}
\partial_s (\gamma_5 X)_1^a \partial_t (\gamma_5 X)_1^b &= \left(-\gamma^\mu (\partial_s \nabla_\mu)_1^a + \frac{1}{2} r (\partial_s \Delta)_1^a \right) \left(\gamma^\nu (\partial_t \nabla_\nu)_1^b + \frac{1}{2} r (\partial_t \Delta)_1^b \right) \\
&= \tilde{V}^{ab} + \text{a term not involving } \gamma^\mu \text{'s} \quad (4.29)
\end{aligned}$$

where

$$\tilde{V}^{ab} = \tilde{V}^{(1)ab} + \tilde{V}^{(2)ab} \quad (4.30)$$

$$\tilde{V}^{(1)ab} = -\frac{1}{2} r \gamma^\mu \left((\partial_s \nabla_\mu)_1^a (\partial_t \Delta)_1^b - (\partial_s \Delta)_1^a (\partial_t \nabla_\mu)_1^b \right) \quad (4.31)$$

$$\tilde{V}^{(2)ab} = -\frac{1}{2} [\gamma^\mu, \gamma^\nu] (\partial_s \nabla_\mu)_1^a (\partial_t \nabla_\nu)_1^b \quad (4.32)$$

It follows from (4.28) and (4.29)–(4.32) that

$$\begin{aligned}\text{Tr}(P[\partial_s P, \partial_t P]) &= \frac{-3}{32} a^4 d^{abc} \text{Tr}(L_0^{-5/2} \gamma_5 X_0 \tilde{V}^{ab} V_2^c) + O(a) \\ &= \frac{-3}{32} a^4 d^{abc} \text{Tr}\left(L_0^{-5/2} \gamma_5 \left(\gamma^\mu \nabla_\mu (\tilde{V}^{(1)ab} V_2^{(2)c} + \tilde{V}^{(2)ab} V_2^{(1)c}) + r \frac{1}{2} \Delta \tilde{V}^{(2)ab} V_2^{(2)c}\right)\right) \\ &\quad + O(a)\end{aligned}\tag{4.33}$$

$V^{(1)ab}$ and $V^{(2)ab}$ can be determined as follows. Recalling $\partial_s U_\mu(x) = a \partial_s A_\mu(x) + O(a^2)$ we get

$$(\partial_s \nabla_\mu^\pm)_1 \psi(x) = (\partial_s A_\mu(x) + O(a)) \psi(x \pm a e_\mu)\tag{4.34}$$

and calculations give

$$\begin{aligned}(\partial_s \nabla_\mu)_1^a (\partial_t \nabla_\nu)_1^b \psi(x) &= \frac{1}{4} (\partial_s A_\mu^a(x) \partial_t A_\nu^b(x) + O(a)) \\ &\quad \times \left[\begin{aligned} &\psi(x + a e_\mu + a e_\nu) + \psi(x + a e_\mu - a e_\nu) \\ &+ \psi(x - a e_\mu + a e_\nu) + \psi(x - a e_\mu - a e_\nu) \end{aligned} \right]\end{aligned}\tag{4.35}$$

$$\begin{aligned}& \left((\partial_s \nabla_\mu)_1^a (\partial_t \Delta)_1^b - (\partial_s \Delta)_1^a (\partial_t \nabla_\mu)_1^b \right) \psi(x) \\ &= - \sum_\nu \frac{1}{2} \left(\partial_s A_\mu^a(x) \partial_t A_\nu^b(x) - \partial_s A_\nu^a(x) \partial_t A_\mu^b(x) + O(a) \right) \\ &\quad \times \left[\begin{aligned} &\psi(x + a e_\mu + a e_\nu) + \psi(x + a e_\mu - a e_\nu) \\ &+ \psi(x - a e_\mu + a e_\nu) + \psi(x - a e_\mu - a e_\nu) \end{aligned} \right]\end{aligned}\tag{4.36}$$

which determine $\tilde{V}^{(1)ab}$ and $\tilde{V}^{(2)ab}$ in (4.31)–(4.32). On the other hand, from (4.11)–(4.12) and (4.13)–(4.16) we find that $V_2^{(1)c}$ and $V_2^{(2)c}$ coincide up to $O(a)$ with $\tilde{V}^{(1)ab}$ and $\tilde{V}^{(2)ab}$, respectively, after replacing $\partial_s A_\mu^a(x) \partial_t A_\nu^b(x)$ with $-\frac{1}{2} F_{\mu\nu}^c(x)$.

Having determined the operators in (4.33) we can now evaluate the trace by first tracing over spinor indices and then evaluating the remaining trace in the plane wave orthonormal basis $\{\phi_k(x)\}$ for periodic scalar fields, given by

$$\phi_k(x) = \frac{1}{\mathcal{N}} e^{ik \cdot x}\tag{4.37}$$

$$\mathcal{N} = (2N)^4, \quad k_\mu \in \frac{\pi}{aN} \{-N, -N+1, \dots, N-1\}\tag{4.38}$$

The result is

$$\text{Tr}(P[\partial_s P, \partial_t P]) = \frac{\mathcal{I}}{32\pi^2} d^{abc} a^4 \sum_x \epsilon_{\mu\nu\rho\sigma} \partial_s A_\mu^a(x) \partial_t A_\nu^b(x) F_{\rho\sigma}^c(x)\tag{4.39}$$

where

$$\mathcal{I} = \sum_k a^4 \Delta^4 k \mathcal{I}_r(ak) \quad (4.40)$$

$$\Delta^4 k \equiv \frac{(2\pi)^4}{a^4 \mathcal{N}} = \text{the “volume per } k \text{” in (4.38),}$$

$$\mathcal{I}_r(k) = \frac{-3r}{8\pi^2} \frac{\prod_{\nu=1}^4 \cos k_\nu \left[-1 + \sum_\mu (1 - \cos k_\mu) - \sum_\mu \frac{\sin^2 k_\mu}{\cos k_\mu} \right]}{\left[\sum_\mu \sin^2 k_\mu + r^2 \left(-1 + \sum_\mu (1 - \cos k_\mu) \right)^2 \right]^{5/2}} \quad (4.41)$$

(The denominator in this expression comes from $L_0^{-5/2}$ in (4.33); we have used (4.17).)

Changing variables from k to ak in (4.40) leads to

$$\lim_{a \rightarrow 0} \mathcal{I} = \int_{-\pi}^{\pi} d^4 k \mathcal{I}_r(k). \quad (4.42)$$

This integral was encountered in [33] in connection with the axial anomaly for fermions with Overlap Dirac operator, and was found to equal 1 (independent of r). We can now take the $a \rightarrow 0$ limit in (4.39) and get the desired result:

$$\lim_{a \rightarrow 0} \text{Tr}(P[\partial_s P, \partial_t P]) = \frac{-1}{32\pi^2} d^{abc} \int_{T^4} d^4 x \epsilon_{\mu\nu\rho\sigma} \partial_s A_\mu^a(x) \partial_t A_\nu^b(x) F_{\rho\sigma}^c(x) \quad (4.43)$$

In deriving this formula we have followed the approach used in [33] for calculating the axial anomaly for the overlap Dirac operator. Presumably the other approaches of ref.'s [34] and [35] could also be used to derive this formula.

Remark. The preceding also shows that

$$\lim_{a \rightarrow 0} \frac{1}{8} \text{tr}\{[\epsilon[\partial_s \epsilon, \partial_t \epsilon]](x, x)\} = \frac{-1}{32\pi^2} d^{abc} \epsilon_{\mu\nu\rho\sigma} \partial_s A_\mu^a(x) \partial_t A_\nu^b(x) F_{\rho\sigma}^c(x) = \frac{-1}{24} \text{tr}(\mathcal{F}(x, s, t))^3 \quad (4.44)$$

Here $[\cdots](x, y)$ denotes the kernel of the operator $[\cdots]$, $\mathcal{F}(x, s, t)$ is the curvature of the gauge field $\mathcal{A}(x, s, t) = A_\mu^{(s,t)}(x) dx^\mu$ in 6 dimensions, and the last coefficient arises as $2\pi i \left(\frac{1}{(2\pi i)^3 3!} \right) = \frac{-1}{24\pi^2}$. This result for the “topological field” $q(x, s, t) = \frac{1}{8} \text{tr}\{[\epsilon[\partial_s \epsilon, \partial_t \epsilon]](x, x)\}$ in 6 dimensions was used in [14]¹⁸ to show the existence of a $w_+(U)$ (or equivalently, the existence of a local gauge-invariant current $j_\mu(x)$ satisfying certain conditions) to all orders in an expansion in the lattice spacing a such

¹⁸ In [14] a more general $q(x, s, t)$ was considered; however, to obtain the mentioned result it suffices to consider the present $q(x, s, t)$.

that the corresponding overlap $\langle v_+, w_+ \rangle$ is gauge-invariant when $d_R^{abc} = 0$. The calculations in this section leading to (4.44) above could therefore be useful as a starting point for finding explicit expressions for the terms in $j_\mu(x)$.

5 Concluding remarks

The main result of this paper is that the overlap formulation of chiral gauge theory on the lattice reproduces the global obstructions to gauge-invariance discussed in the continuum by Alvarez-Gaumé and Ginsparg [3]. We showed that the obstruction on the lattice reduces to the continuum obstruction in the classical continuum limit. This, together with the fact that the lattice obstruction is also an integer (winding number), implies that the lattice obstruction coincides exactly with the continuum one for small non-zero lattice spacing (i.e. close to the classical continuum limit). Thus the overlap formulation is seen to exactly capture topological structure of the continuum theory in the nonabelian case, just as it does in the abelian case considered previously in [5]. We mention again that, while we have taken the spacetime to be the 4-dimensional, our arguments and results generalise straightforwardly to Euclidean spacetime T^{2n} for arbitrary n .

It might be instructive to compare this with the situation for chiral Wilson fermions on the lattice (where gauge-invariance is explicitly broken due to the Wilson term, and only restored in the $a \rightarrow 0$ limit). In this case the consistent local anomaly $\frac{d}{d\theta} \log \det(D_{Wilson}^{\phi_\theta \cdot U})_+$ has been shown to converge to the continuum anomaly in the classical continuum limit [36], so the integral

$$W_{Wilson}(\Phi) = \frac{1}{2\pi i} \int_0^1 d\theta \frac{d}{d\theta} \log \det(D_{Wilson}^{\phi_\theta \cdot U})_+$$

converges to the continuum global obstruction $\deg(\Phi)$ in this limit. However, for non-zero lattice spacing the integral $W_{Wilson}(\Phi)$ is *non-integer* in general –it does not have a winding number interpretation since $|\det(D_{Wilson}^U)_+|$ is not gauge-invariant. This is in contrast to the overlap case where

$$W(\Phi) = \frac{1}{2\pi i} \int_0^1 d\theta \frac{d}{d\theta} \log \langle v_+, w_+(\phi_\theta \cdot U) \rangle$$

is an integer (winding number) since $|\langle v_+, w_+(U) \rangle|$ is gauge-invariant and $\langle v_+, w_+(\phi_\theta \cdot U) \rangle$ differs from $\langle v_+, w_+(U) \rangle$ only by a phase factor. Thus, in contrast to the overlap case, the non-integer $W_{Wilson}(\Phi)$ is in general not equal to the integer-valued continuum obstruction $\deg(\Phi)$ at any non-zero lattice spacing; equality only occurs in the limit $a \rightarrow 0$.

As a consequence of our main result, we found that the continuum anomaly cancellation condition $d_R^{abc} = 0$ is a necessary condition for anomaly cancellation in the overlap on the lattice (at least when $Map(T^5, SU(N))$ contains maps with non-trivial degree). While this is no surprise, our derivation is robust compared to other approaches: Firstly, it is independent of the choice of phase in the overlap (in contrast to the consistent local anomaly which does depend on the phase choice), and secondly, no approximations, or assumptions on the gauge field, have been used.

In the continuum argument of ref. [3], the winding number obstruction $W(\Phi)$ is shown to equal the index of a Dirac operator $\mathcal{D}$ in $6 = 2n + 2$ dimensions. The index theorem is then used to get $\text{index } \mathcal{D} = \deg \Phi$. In this paper we have followed a different route: a lattice version of the determinant line bundle approach of ref. [4]. In fact, it is also possible to give a lattice version of the original argument of Alvarez-Gaumé and Ginsparg in the overlap setting, using a certain lattice Dirac operator in $2n + 2$ dimensions (with the extra 2 dimensions being continuous) [37]¹⁹

The obstructions of Alvarez-Gaumé and Ginsparg are but one type of obstruction to gauge-invariance of the chiral determinant. In general, the obstructions are manifestations of non-trivial topological structure of the determinant line bundle over the orbit space of gauge fields. This topic has been studied in detail in the continuum; see, e.g., [4]. The results of [5] and the present paper suggest that in general the continuum topological structure of the determinant line bundle can be reproduced on the lattice in the overlap formulation (at least when the spacetime manifold is a $2n$ -dimensional torus). The determinant line bundle comes equipped with a canonical

¹⁹In this case, the index density of the lattice $\mathcal{D}$ turns out to be the aforementioned topological field $q(x, s, t)$ in $6 = 2n + 2$ dimensions which appeared in ref. [14].

$U(1)$ connection, and the difference $Im\Gamma(A^{(1)}) - Im\Gamma(A^{(0)})$ for the effective action $\Gamma(A) = \log \det(D_+^A)$ can be expressed in terms of the parallel transport of this connection along a path joining $A^{(0)}$ to $A^{(1)}$ [38]. This can in turn be expressed in terms of a spectral flow (η -invariant) of a Dirac operator and a Chern-Simons term, both in $5 = 2n + 1$ dimensions. Lattice versions of these relations in the overlap setting have already been found [11, 28].

Finally, it is interesting to note that the quantity $\text{Tr}(PdPdP) = \frac{1}{8}\text{Tr}(\epsilon d\epsilon d\epsilon)$, which in the present setting appears as the curvature of the overlap determinant line bundle (or the 'Berry curvature' in the terminology of ref. [5]), also arises as the curvature of a determinant line bundle in canonical quantisation of the continuum theory.²⁰ See [39] and the ref.'s therein. In that setting one considers a certain infinite-dimensional Grassmannian manifold consisting of splittings $V_+ \oplus V_-$ of the Hilbert space of 1-particle states; each splitting corresponds to an $\epsilon = P_{V_+} - P_{V_-}$. There is a canonical determinant line bundle on this manifold, and its curvature turns out to be a renormalised version of $\frac{1}{8}\text{Tr}(\epsilon d\epsilon d\epsilon)$.²¹ It could be interesting to explore the apparent analogy between this continuum formulation and the lattice overlap formulation. Recently, an obstruction to canonical quantisation of the continuum theory on odd-dimensional spacetimes was described in [41]. Instead of the 'Berry curvature' 2-cocycle, the obstruction there is given in terms of a 3-cocycle known as the Dixmier–Douady class. It could be interesting to see if there is something analogous to this in the lattice overlap formulation.

Acknowledgements. This work was presented at the mini-workshop “New developments in lattice gauge theory”, CSSM, Adelaide, 4-5 April'00, and I thank Prof.'s Ting-Wai Chiu, Kazuo Fujikawa, Urs Heller, Jouko Mickelsson and Tony Williams for interesting comments and discussions at that time. I also thank Prof. Her-

²⁰ I thank Prof. J. Mickelsson for pointing this out to me.

²¹ In fact, a lattice regularisation of chiral gauge theory (different to the overlap) in which this quantity also appears has been presented in ref. [40]. Gauge-invariance appears to be problematic in this approach though: the notion of gauge symmetry needs to be modified on the lattice in a way that involves non-local operators.

bert Neuberger for correspondence on the Overlap and, together with Prof.'s Martin Lüscher and Jun Nishimura, for feedback on earlier versions of this paper. The author is supported by an ARC postdoctoral fellowship.

Appendix

A Derivation of (3.7): $\frac{\langle v_+, (dw_+)_{\perp} \rangle}{\langle v_+, w_+ \rangle} = \text{Tr}(dDD^{-1}P_+)$.

Let $U(t)$ be a smooth curve in $\mathcal{U}$. Using (2.13) we calculate

$$\begin{aligned} \frac{d}{dt} \log \langle v_+, w_+ \rangle &= \frac{d}{dt} \log \langle v_+, \widehat{D}w_+ \rangle \\ &= \frac{1}{\langle v_+, \widehat{D}w_+ \rangle} \left(\langle v_+, \frac{d}{dt} \widehat{D}w_+ \rangle + \langle v_+, \widehat{D} \frac{d}{dt} w_+ \rangle \right) \end{aligned} \quad (\text{A.1})$$

At $t = 0$ (and with our assumption that D has no zero-modes) we have

$$\begin{aligned} \langle v_+, \frac{d}{dt} \widehat{D}w_+ \rangle &= \langle v_+, \frac{d}{dt} \widehat{D} \widehat{D}^{-1} \widehat{D}w_+ \rangle = \langle v_+, \widehat{D}w_+ \rangle \langle v_+, \frac{d}{dt} \widehat{D} \widehat{D}^{-1} v_+ \rangle \\ &= \langle v_+, \widehat{D}w_+ \rangle \frac{d}{dt} \Big|_{t=0} \det \left(D(t) D(0)^{-1} \Big|_{\mathcal{C}_+} \right) \\ &= \langle v_+, \widehat{D}w_+ \rangle \text{Tr} \left(\frac{d}{dt} D D^{-1} P_+ \right) \end{aligned}$$

so the first term on the right-hand side of (A.1) is $\text{Tr}(\frac{d}{dt} D D^{-1} P_+)$. Comparing (A.1) with (3.6) we see that (3.7) holds iff

$$\langle v_+, \widehat{D} \frac{d}{dt} w_+ \rangle = \langle v_+, \widehat{D}w_+ \rangle \langle w_+, \frac{d}{dt} w_+ \rangle \quad (\text{A.2})$$

Choose an orthonormal basis $w_1(t), \dots, w_d(t)$ for $\mathcal{C}_+^{(1, U(t))}$ such that $w_+ = w_1 \wedge \dots \wedge w_d$. Then

$$\langle v_+, \widehat{D} \frac{d}{dt} w_+ \rangle = \sum_{j=1}^d \langle v_+, \widehat{D}(w_1 \wedge \dots \wedge \frac{d}{dt} w_j \wedge \dots \wedge w_d) \rangle \quad (\text{A.3})$$

Substitute $\frac{d}{dt} w_j = \sum_{k=1}^d \langle w_k, \frac{d}{dt} w_j \rangle w_k + (\frac{d}{dt} w_j)_{\perp}$ in (A.3), where $(\frac{d}{dt} w_j)_{\perp} \in \mathcal{C}_-^{(1, U(t))}$. Since D maps $\widehat{\mathcal{C}}_{\pm} = \mathcal{C}_{\mp}^{(1, U)}$ to $\mathcal{C}_{\mp}$, the terms involving $(\frac{d}{dt} w_j)_{\perp}$ give vanishing contribution, and we get

$$\langle v_+, \widehat{D} \frac{d}{dt} w_+ \rangle = \sum_{j=1}^d \langle v_+, \widehat{D}w_+ \rangle \langle w_j, \frac{d}{dt} w_j \rangle.$$

This equals the right-hand side of (A.2) as required, since

$$\begin{aligned}\langle w_+, \frac{d}{dt} w_+ \rangle &= \sum_{j=1}^d \langle w_+, w_1 \wedge \cdots \wedge \frac{d}{dt} w_j \wedge \cdots \wedge w_d \rangle \\ &= \sum_{j=1}^d \langle w_+, w_+ \rangle \langle w_j, \frac{d}{dt} w_j \rangle = \sum_{j=1}^d \langle w_j, \frac{d}{dt} w_j \rangle.\end{aligned}$$

B Derivation of (3.14)

It suffices to restrict to a surface in $\mathcal{U}$ with coordinates (s, t) and show

$$\langle \partial_s w_+, \partial_t w_+ \rangle - \langle \partial_t w_+, \partial_s w_+ \rangle = \text{Tr}(P \partial_s P \partial_t P) - \text{Tr}(P \partial_t P \partial_s P) \quad (\text{B.1})$$

Let $w_1(s, t), \dots, w_d(s, t)$ be an orthonormal basis for $\mathcal{C}_+^{(1, U(s, t))}$ such that $w_+ = w_1 \wedge \cdots \wedge w_d$. It is convenient to use bra-ket notation:

$$P = \sum_{k=1}^d |w_k\rangle \langle w_k|, \quad \partial_s P = \sum_{k=1}^d |\partial_s w_k\rangle \langle w_k| + |w_k\rangle \langle \partial_s w_k|$$

Then

$$\begin{aligned}\text{Tr}(P \partial_s P \partial_t P) &= \sum_{j=1}^d \langle w_j | \partial_s P \partial_t P | w_j \rangle \\ &= \sum_{j,k,l=1}^d \langle w_j | \left(|\partial_s w_k\rangle \langle w_k| + |w_k\rangle \langle \partial_s w_k| \right) \left(|\partial_t w_l\rangle \langle w_l| + |w_l\rangle \langle \partial_t w_l| \right) | w_j \rangle \\ &= \sum_{j,k} \langle w_j | \partial_s w_k \rangle \langle w_k | \partial_t w_j \rangle + \sum_{j,k} \langle w_j | \partial_s w_k \rangle \langle \partial_t w_k | w_j \rangle \\ &\quad + \sum_j \langle \partial_s w_j | \partial_t w_j \rangle + \sum_{k,l} \langle \partial_s w_j | w_l \rangle \langle \partial_t w_l | w_j \rangle\end{aligned} \quad (\text{B.2})$$

The first and fourth sums are clearly symmetric under $\partial_s \leftrightarrow \partial_t$. The second sum is likewise symmetric under $\partial_s \leftrightarrow \partial_t$ as is easily seen using $\langle \delta w_j | w_k \rangle + \langle w_j | \delta w_k \rangle = \delta \langle w_j | w_k \rangle = 0$. It follows that

$$\text{Tr}(P \partial_s P \partial_t P) - \text{Tr}(P \partial_t P \partial_s P) = \sum_j \langle \partial_s w_j | \partial_t w_j \rangle - \langle \partial_t w_j | \partial_s w_j \rangle$$

and this is equal to the left-hand side of (B.1) as required.

References

- [1] R. Narayanan and H. Neuberger, Phys. Lett. B **302** (1993) 62; Phys. Rev. Lett **71** (1993) 3251; Nucl. Phys. B **412** (1994) 574
- [2] R. Narayanan and H. Neuberger, Nucl. Phys. B **443** (1995) 305
- [3] L. Alvarez-Gaumé and P. Ginsparg, Nucl. Phys. B **243** (1984) 449
- [4] M. F. Atiyah and I. M. Singer, Proc. Natl. Acad. Sci. USA **81** (1984) 2597; J.-M. Bismut and D. S. Freed, Commun. Math. Phys. **106** (1986) 159; **107** (1986) 103.
- [5] H. Neuberger, Phys. Rev. D **59** (1999) 085006
- [6] R. Narayanan and J. Nishimura, Nucl. Phys. B **508** (1997) 371
- [7] Y. Kikukawa and H. Neuberger, Nucl. Phys. B **513** (1998) 735
- [8] H. Neuberger, Phys. Lett. B **437** (1998) 117
- [9] O. Bär and I. Campos, Nucl. Phys. B **581** (2000) 499
- [10] S. Randjbar-Daemi and J. Strathdee, Phys. Lett B **348** (1995) 543; Nucl. Phys. B **443** (1995) 386; **466** (1996) 335; Phys. Lett. B **402** (1997) 134
- [11] H. Suzuki, Prog. Theor. Phys. **101** (1999) 1147
- [12] H. Suzuki, hep-lat/0002009
- [13] M. Lüscher, Nucl. Phys. B **549**, 295 (1999)
- [14] M. Lüscher, Nucl. Phys. B **568** (2000) 162
- [15] H. Neuberger, hep-lat/0002032
- [16] M. Golterman and Y. Shamir, Nucl. Phys. Suppl. **47** (1996) 603
- [17] R. Narayanan and H. Neuberger, Phys. Lett. B **358** (1995) 303;

- [18] T. Izubuchi and J. Nishimura, JHEP **9910:002** (1999)
- [19] R. Narayanan and H. Neuberger, Nucl. Phys. B **477** (1996) 521; Phys. Lett. B **402** (1997) 320; Y. Kikukawa, R. Narayanan and H. Neuberger, Phys. Lett B **399** (1997) 105; Phys. Rev. D **57** (1998) 1233
- [20] Y. Kikukawa, Nucl. Phys. (Proc. Suppl.) **63** (1998) 587
- [21] P. Ginsparg and K. G. Wilson, Phys. Rev. D **25** (1982) 2649
- [22] P. Hasenfratz and F. Niedermayer, Nucl. Phys. B **414** (1994) 785; P. Hasenfratz, in Non-perturbative quantum field physics (Peñíscola 1997), eds. M. Asorey, A. Dobado (World Scientific, Singapore, 1998); Nucl. Phys. B (Proc. Suppl.) **63A–C** (1998) 53; Nucl. Phys. B **525** (1998) 401.
- [23] P. Hasenfratz, V. Laliena and F. Niedermayer, Phys. Lett. B **427** (1998) 125
- [24] M. Lüscher, Nucl. Phys. B **538**, 515 (1999)
- [25] T. Fujiwara, H. Suzuki and K. Wu, Phys. Lett. B **463** (1999) 63; Nucl. Phys. B **569** (2000) 643; hep-lat/9910030; hep-lat/0001029; H. Suzuki, hep-lat/99110009
- [26] P. Hernández, K. Jansen and M. Lüscher, Nucl. Phys. B **552** (1999) 363
- [27] H. Neuberger, Phys. Rev. D **61** (2000) 085015
- [28] T. Aoyama and Y. Kikukawa, hep-lat/9905003
- [29] H. Neuberger, hep-lat/9911022; hep-lat/9912013
- [30] H. Neuberger, Phys. Lett. B **417** (1998) 141; **427** (1998) 353
- [31] J. E. Avron, R. Seiler, B. Simon, Phys. Rev. Lett. **51** (1983) 51
- [32] D. H. Adams, in preparation
- [33] D. H. Adams, hep-lat/9812003; H. Suzuki, Prog. Theor. Phys. **102** (1999) 141

- [34] Y. Kikukawa and A. Yamada, Phys. Lett. B **448** (1999) 265
- [35] K. Fujikawa, Nucl. Phys. B **546** (1999) 480
- [36] S. Aoki, Phys. Rev. D **35** (1986) 1435; A. Coste, C. Korthals Altes and O. Napoly, Nucl. Phys. B **289** (1987) 645; T. Jolicoeur, R. Lacaze and O. Napoly, Nucl. Phys. B **293** (1987) 215
- [37] D. H. Adams, hep-lat/9910036 v2
- [38] S. Della Pietra, V. Della Pietra and L. Alvarez-Gaumé, Commun. Math. Phys. **109** (1987) 691
- [39] J. Mickelsson, Commun. Math. Phys. **154** (1993) 403
- [40] J. Mickelsson, hep-lat/9602030
- [41] A. L. Carey and J. Mickelsson, hep-th/9912003